

- Heufelder, George R.; Schneeberger, Philip J.; Madenjian, Charles P.; Rutecki, Thomas L.; Mansfield, Pamela J.; Auer, Nancy A.; Noguchi, George E. (1981): Special Report Number 86 Great Lakes Research Division, The University of Michigan: Adult, Juvenile, and Larval Fish Populations in the Vicinity of the J.H. Campbell Power Plant, Eastern Lake Michigan, 1977-1980.
- Heufelder, G.R.; Jude, D.J.; and Tesar, F.J. (1982): Effects of Upwelling on Local Abundance and Distribution of Larval Alewife (*Alosa pseudoharengus*) in Eastern Lake Michigan, Volume 39, Number 11, pgs. 1531-1537.
- Hunsaker, C.T.; Carpenter, D.E. (1990): Environmental Monitoring and Assessment Program: Ecological Indicators. U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC. [INDICATOR, PROGRAM]
- Illinois Environmental Protection Agency (): Lake Michigan Water Quality Report 1985. 174 Pages. [GREAT LAKES, LAKE MICHIGAN, WATER QUALITY, TOXICS, PLANKTON, CHLORIDE]
- Illinois Environmental Protection Agency (1988): Lake Michigan Water Quality Report 1986. 128 Pages. (IEPA/WPC/88-015). [GREAT LAKES, LAKE MICHIGAN, WATER QUALITY, TOXICS, PLANKTON, CHLORIDE]
- Illinois State Environmental Protection Agency, Springfield, Div. of Water Pollution Control (1988): Lake Michigan Water Quality Report, 1986. 140 pages. Section 305(b) Report (EPAIWPC/88-015) [EUTROPHICATION, FISH, HUMAN USE, LAKE MICHIGAN, POLLUTION, TOXICS, WASTE DISPOSAL, WATER COLUMN]
- Illinois Environmental Protection Agency (1992): Illinois Water Quality Report 1990-1991. Section 305 (b) Report. 405 pages. [EUTROPHICATION, FISH, HABITAT, HUMAN HEALTH, HUMAN USE, SEDIMENT, TOXICS, WATER COLUMN, WETLAND]
- Illinois Environmental Protection Agency, Bureau of Water, Division of Water Pollution Control (1993): Lake Michigan Water Quality Report 1989 - 1991. 135 Pages. (IEPA/WPC/93-065) [HEALTH, FISH, GREAT LAKES, HUMAN HEALTH, LAKE MICHIGAN, LEAD, PCB, PESTICIDE, PLANKTON, POLLUTION, RECREATION, REGULATION WATER QUALITY, WATER QUALITY STANDARDS]
- Illinois Environmental Protection Agency, Bureau of Water (1993): Water Quality in Illinois (1990-1991). 9 Pages. (IEPA/WPC/92-224). [WATER QUALITY, GREAT LAKES, RIVERS, STREAMS, FISH, SEDIMENT, TOXICS]
- Indiana Department of Environmental Management (1991): The Remedial Action Plan for the Indiana Harbor Canal, The Grand Calumet River, and the Nearshore Lake Michigan - Stage One. 84 Pages. [LAKE MICHIGAN, WATER QUALITY]

Indiana State Board of Health, Division of Water Pollution Control (): Physical & Chemical Characteristics of the Indiana Portion of Lake Michigan 1980-1981. Prepared for the Indiana Environmental Management Board, Indiana Stream Pollution Control Board. 76 Pages. [GREAT LAKES, LAKE MICHIGAN, WATER QUALITY, CHLORIDES]

International Joint Commission (1982): A Review of the Pollution Abatement Programs Relating to the Petroleum Refinery in the Great Lakes Basin. [REFINERY, POLLUTION ABATEMENT, CANADA, US, EFFLUENT LIMITS]

International Joint Commission (1983): An Inventory of Chemical Substances Identified in the Great Lakes Ecosystem. Vol. 1-6. International Joint Commission, Great Lakes Water Quality Board, Windsor, Ontario. [ECOSYSTEM, GREAT LAKES, TOXICS]

International Joint Commission Surveillance Work Group (1985): Guidance on Characterization of Toxic Substances Problems in Areas of Concern in the Great Lakes Basin (1985). [TOXICS, TOXICS PROBLEMS IN THE GREAT LAKES]

International Joint Commission (1986): Great Lakes International Surveillance Plan (2 Vols)., Windsor, Ontario. [GREAT LAKES]

International Joint Commission (1986): Report of the Aquatic Ecosystem Objectives Committee, 1985 Annual Report Great Lakes Science Advisory Board Report to the International Joint Commission. [ECOSYSTEM]

International Joint Commission, Great Lakes Regional Office. (1988). Report on Modeling the Loading-Concentration Relationship for Critical Pollutants in the Great Lakes, Prepared by the Task Force on Chemical Loadings of the Toxic Substances Committee, October. [GREAT LAKES, TOXICS, LOADINGS]

International Joint Commission (1989): Fifth Biennial Report on Great Lakes Water Quality Part II Under the Great Lakes Water Quality Agreement of 1978 to the Governments of the United States and Canada and the State and Provincial Governments of the Great Lakes Basin. 57 pages. [EXOTIC SPECIES, FISH, GREAT LAKES, HUMAN HEALTH, LAKE SUPERIOR, PROGRAM, TOXICS, WATER COLUMN]

International Joint Commission (1989): 1989 Report on Great Lakes Water Quality. Great Lakes Water Quality Board. [WATER COLUMN]

International Joint Commission, United States and Canada (1990): Fifth Biennial Report on Great Lakes Water Quality. International Joint Commission, Washington, DC. (ISBN 1-895085-03-09) [GREAT LAKES, WATER COLUMN]

International Joint Commission and Great Lakes Fishery Commission (1990); Exotic Species and the Shipping Industry: The Great Lakes-St. Lawrence Ecosystem at Risk. A special report to the Governments of the United States and Canada. 77 pages. [ECOSYSTEM, EXOTIC SPECIES, GREAT LAKES]

International Joint Commission, Virtual Elimination Task Force (1991): Persistent Toxic Substances: Virtually Eliminating Inputs to the Great Lakes. Interim report., Windsor, Ontario. 42 pages. [GREAT LAKES, LOADINGS, TOXICS, WATER COLUMN]

International Joint Commission (1992): Sixth Biennial Report on Great Lakes Water Quality Under the Great Lakes Water Quality Agreement of 1978 to the Governments of the United States and Canada and the State and Provincial Governments of the Great Lakes Basin. 60 pages. [ECOSYSTEM, GREAT LAKES, LAKE SUPERIOR, PROGRAM; RESTORATION; TOXICS, WATER COLUMN]

International Joint Commission: Seventh Biennial Report on Great Lakes Water Quality. [WATER QUALITY, GREAT LAKES]

International Joint Commission (1993). Great Lakes Science Advisory Board. Report to the International Joint Commission.

International Joint Commission (1994): Bibliography of Reports Issued Under the Boundary Waters Treaty of 1909 and Great Lakes Water Quality Agreements of 1972 and 1978, and the Protocol Amending the 1978 Agreement.

International Joint Commission Reference Group; International Reference Group on Great Lakes Pollution (From Land Use Activities) Trophic Characterization of the US and Canadian Nearshore Zones of the Great Lakes. [TROPIC LEVELS, TROPIC STATUS, NEARSHORE ZONES, CHLOROPHYLL, PHOSPHOROUS, EUTROPHICATION]

International Joint Commission. Proceedings of the Roundtable on Contaminant-caused Reproductive Effects in Salmonids. [REPRODUCTIVE EFFECTS, SALMON, SALMONIDS, TROUT, CHLORINATED ORGANICS]

Jacobson, Joseph L.; Jacobson, Sandra W. (1993): A 4-Year Followup Study of Children Born to Consumers of Lake Michigan Fish. J. Great Lakes Res. 19(4), Pages 776-783. [GREAT LAKES, FISH, HEALTH]

Janssen, John; and Quinn, John. (19__): Chapter 17 - Biota of the Naturally Rocky Area of Southwestern Lake Michigan with Emphasis on Potential Fish Prey.

Jude, David J: Entrainment of Fish Larvae and Eggs on the Great Lakes, With Special Reference to the D.C. Cook Nuclear Plant, Southeastern Lake Michigan. [FISH, PLANKTON, LAKE MICHIGAN]

Jude, D.J.; Klinger, S.A.; Enk, M.D. (1981): First Evidence of Natural Reproduction by Planted Lake Trout in Lake Michigan. Journal of Great Lakes Research 7, 57-61. [FISH, LAKE MICHIGAN, REPRODUCTION]

Jude, David J.; and Tesar, Frank J. (1985): Recent Changes in the Inshore Forage Fish of Lake Michigan. Canadian Journal of Fisheries and Aquatic Sciences, Volume 42, Number 6, Pp. 1154-1157. [FISH, LAKE MICHIGAN]

- Jude, D.J.; Heang T.T.; Heufelder, G.R.; et.al. : Adult, Juvenile and Larval Fish Populations in the Vicinity of the J.H. Campbell Power Plant, Eastern Lake Michigan, 1977-1980. Special Report No. 86 of the Great Lakes Research Division. University of Michigan.
- Jude, David J.; Reider, Robert H.; Smith, Gerald R. (1992): Establishment of Gobiidae in the Great Lakes Basin. *Canadian Journal of Fisheries and Aquatic Sciences* 49(2), 416-421. [EXOTIC SPECIES, GREAT LAKES]
- Jude, D.J. (1992): Evidence for Natural Reproduction by Stocked Walleyes in the Saginaw River Tributary System, Michigan. *North American Journal of Fisheries Management* 12:386-395. [FISH, GREAT LAKES]
- Jude, David J.; Pappas, Janice (1992): Fish Utilization of Great Lakes Coastal Wetlands. *Journal of Great Lakes Research* 18, 651-672. [FISH, GREAT LAKES, HABITATS WETLAND]
- Jude, D.J.; Leach, J. (1993): The Great Lakes Fisheries. Chap. 22. In: *North American Fisheries Management Textbook of American Fisheries Society*. (Eds: Kohler, C; Hubert, W), Bethesda, MD. [FISH, GREAT LAKES]
- Keeler, Gerald J.; Pacyna, Jezef M.; Bidleman, Terry F.; Nriagu, Jerome O. (1993): Identification of Sources Contributing to the Contamination of the Great Waters by Toxic Compounds. 145 pages. [LOADINGS, TOXICS]
- Kelly, J.R.; Harwell, M.A. (1989): Indicators of ecosystem response and recovery. In: *Ecotoxicology: Problems and Approaches*. (Eds: Levin, S.A.; Harwell, M.A.; Kelly, J.R., Kimball, K.D.) Springer Verlag, New York, NY, 9-40. [ECOSYSTEM, INDICATOR, TRENDS]
- Kenaga, D.E., Creal, W.S., and Basch, R.E.: *Limnology of Michigan's Nearshore Waters of Lake Michigan*. U.S. EPA: 100 pp. EPA 905/3-83-003.
- Kielty, T.J.; Landrum, P.F. (1990): Population-Specific Toxicity Responses by the Freshwater Oligochaete *Stylocrilius heringianus*, in Natural Lake Michigan Sediments. *Environmental Toxicology and Chemistry* 9, 1147-1154. [BENTHOS, LAKE MICHIGAN, SEDIMENT]
- Kitchell, J.F.; Evans, M.S.; Scavia, D.; Crowder, L.B. (1988): Regulation of water quality in Lake Michigan: Report of the Food Web Workshop. *Journal of Great Lakes Research* 14, 109-114. [EUTROPHICATION, LAKE MICHIGAN, PROGRAM, WATER COLUMN]
- Koonce, Joseph F. (1990): Commentary on Fish Community Health: Monitoring and Assessment in Large Lakes. *Journal of Great Lakes Research* 16, 631-634. [ECOSYSTEM, FISH, GREAT LAKES, HEALTH]
- Kratt, C.E. (1993): Changes in the Fish Community of Lower Green Bay, Lake Michigan. In: *IAGLR Conference, Green Bay, Wisconsin, June 1993*. [FISH, GREEN BAY]
- Krueger, C.C.; May, B. (1991): Ecological and Genetic Effects of Salmonid Introductions in North America. *Canadian Journal of Fisheries and Aquatic Sciences* 44 (Suppl. 1), 66-77. [FISH]

- Leatherland, John F. (1993): Field Observations on Reproductive and Development Dysfunction in Introduced and Native Salmonids from the Great Lakes. J. Great Lakes Res. 19(4), Pages 737-751. [GREAT LAKES, TOXICS]
- Lee, D.; Eedford, K.; Yen, C. (1993): Storm and Entrainment Effects on Tributary Sediment Loads. J. Hydraulic Engrg. 119(12). [LOADINGS, SEDIMENT]
- Leonard, D.; Manam, R. (1988): Reassessing Estimates of Great Lakes Consumptive Use and Impacts. In: The Great Lakes: Living with North America's Inland Waters. (Ed: Kickcox, D) American Water Resources Association Tech. Pub TPS48-3, Bethesda, MD, 33-44. [GREAT LAKES, HUMAN USE]
- Lesht, B.M.: Michigan Eutrophication Model: Calibration, Sensitivity, and 5-Year Hindcast Analysis. U.S. EPA/Argonne National Laboratory: September 1984. 55 pp.
- Lesht, B.M., and Rockwell, D.C.: The State of the Middle Great Lakes: Results of the 1983 Water Quality Survey of Lakes Erie, Huron, and Michigan. U.S. EPA August 1985, 171 pp. (ANL/ER-84-3).
- Lesht, B.M.; Fontaine, T.D. III; Dolan, D.M. (1991): Great Lakes Total Phosphorus Model: Post Audit and Regionalized Sensitivity Analysis. Journal of Great Lakes Research 17, 3-17. [EUTROPHICATION, GREAT LAKES, PHOSPHORUS, WATER COLUMN]
- Likens, G.E., ED. (1972): Nutrients and Eutrophication: The Limiting Nutrient Controversy Proceedings of the Symposium on Nutrients and Eutrophication W.K. Kellogg Biological Station, Michigan State U. 11 and 12 February 1971. American Society of Limnology and Oceanography, Inc.
- Locke, A; Reid, D.M.; Sprules, W.G.; Carlton, I.T.; van Leeuwen, J.C. (1991): Effectiveness, of Mid-Ocean Exchange in Controlling Freshwater and Coastal Zooplankton in Ballast Water. Can. Tech. Rep. Fish. Aquat Sci. No. 1822. [PLANKTON]
- Loucks, O.L. (1982): Concern for Acidic Deposition in the Great Lakes Region Chap. 2. In: Acid Precipitation: Effects on the Ecological Systems. 21-41. (ISBN-0-250-10509, EPA/600/D-89/239 Grant R-8094 12). [ATMOSPHERE, GREAT LAKES, POLLUTION, WATER COLUMN]
- LTI, Limno-Tech, Inc. (1993): Great Lakes Environmental Assessment. Prepared for: The National Council of the Paper Industry for Air and Stream Improvement (NCASI). [SEDIMENTS, FISH, TOXICS, GREAT LAKES, LAKE MICHIGAN, HEALTH, HABITAT]
- Lyons, J.; Bedford, K.; Yen, C.; Lee, D.; Mark, D. (1988): Determination of Suspended Concentrations from Multiple Day Landsat and AVHRR data. Journal of Remote Sensing of the Environment 24, 364-373. [SEDIMENT]

- Mac, Michael; Gilbertson, Michael (Eds.) (1990): Great Lakes Science Advisory Board's Biological Effects Subcommittee of the Ecological Committee. Report to the International Joint Commission. Proceedings of the Roundtable on Contaminant-Caused Reproductive Problems in Salmonids., Windsor, Ontario. 45 pages. [DEFORMITIES, FISH, GREAT LAKES, LAKE ERIE, LAKE MICHIGAN, LAKE ONTARIO]
- Mac, Michael J.; Edsall, Carol C. (1991): Environmental Contamination and the Reproductive Success of Lake Trout in the Great Lakes an Epidemiological Approach. Journal of Toxicology and Environmental Health 33, 375-394. [EPIDEMIOLOGY, FISH, GREAT LAKES, PCB, POPULATION, REPRODUCTION, TOXICS]
- Mac, Michael J.; Schwartz, Ted R.; Edsall, Carol C.; Frank, Anthony M. (1993): Polychlorinated Biphenyls in Great Lakes Lake Trout and Their Eggs: Relations to Survival and Congener Composition 1979-1988. J. Great Lakes Res. 19(4), Pages 752-765. [FISH, GREAT LAKES]
- Mackay, D.; Paterson, S. (1981): Calculating Fugacity. Environmental Science and Technology 15, 1006-1014. [ECOSYSTEM]
- MacKay, Donald (1990): Modeling pathways for toxic chemicals. In: Great Lakes Monograph No.3. Contemporary and Emerging Issues in the Great Lakes. Proceedings from a Colloquium Between The University at Buffalo and the University of Toronto April 12, 1989. (Eds: Bankert, Lynne S; Flint, R Warren) Great Lakes Program. State University of New York, Buffalo, New York, 33-35. [LAKE ONTARIO, PCB, TOXICS]
- Madenjian, C.P.; Jude, D.J. (1985): Comparison of Sleds Versus Plankton Nets for Sampling Fish Larvae and Eggs. Hydrobiologia 124, Pages 275-281. [FISH, GREAT LAKES]
- Magnuson, John J. (1992): A Sense of Beginning the Laurentian Great Lakes in the Process of Rehabilitation. Restor. Manag. Notes 10, 29-32. [ECOSYSTEM, GREAT LAKES, RESTORATION]
- Maguire, L. (1988): Decision analysis: An integrated approach to ecosystem exploitation and rehabilitation. In: Rehabilitating Damaged Ecosystems. (Ed: Cairns, Jr, J) CRC Press, Boca Raton, FL, 105-122. [ECOSYSTEM; PROGRAM]
- Makarewicz, J.C. (1987). Phytoplankton and Zooplankton Composition, Abundance, and Distribution: Lake Erie, Lake Huron, and Lake Michigan-1983. Vol. 1: Interpretive Report. GLNPO Report 87-05. USEPA: July 1987, 280 p. (EPA-905/3-84-006). [AQUATIC LIFE: FISH STUDIES AND OTHER BIOTA]
- Makarewicz, J.C. (1987): Phytoplankton and Zooplankton in Lakes Erie, Huron and Michigan: 1984. Volume 1, Interpretive Report, Final rept 1984-1985. Environmental Protection Agency, Chicago, IL. Great Lakes National Program Office. 295 pages. (EPA/905/348/001, GLPNO-88/03 Grant EPA-R-005772) [EUTROPHICATION, FISH, GREAT LAKES, INDICATOR, LAKE ERIE, LAKE HURON, LAKE MICHIGAN, NUTRIENTS, OPEN WATER, PLANKTON; POLLUTION, TRENDS, WATER COLUMN]

- Makarewicz, J.C.; Bertram, P.E.; Lewis, T. (1989): Phytoplankton and Zooplankton in Lakes Erie, Huron, and Michigan:1985. Volume 1. Interpretive Report. Final Rept. Environmental Protection Agency, Chicago, IL Great Lakes National Program Office. 272 pages. (EPA/905/3-90/003, GLNPO- 01/91) [ECOSYSTEM; EUTROPHICATION, GREAT LAKES, INDICATOR, LAKE ERIE, LAKE HURON, LAKE MICHIGAN, PLANKTON, POLLUTION, WATER COLUMN]
- Makarewicz, Joseph C.; Bertram, Paul (1991): A lakewide comparison study of phytoplankton biomass and its species composition in Lake Huron, 1971 to 1985. *Journal of Great Lakes Research* 17, 553- 564. [LAKE HURON, OPEN WATER, PLANKTON, TRENDS, WILDLIFE]
- Makarewicz, Joseph C. (1991): Photosynthetic parameters as indicators of ecosystem health. *Journal of Great Lakes Research*. 17, 333-343. [INDICATOR, LAKE ONTARIO]
- Makarewicz, Joseph C. (1993): Phytoplankton Biomass and Species Composition In Lake Erie, 1970 to 1987. *Journal of Great Lakes Research* 19, 258-272. [EUTROPHICATION, LAKE ERIE, PLANKTON, WATER COLUMN]
- Mandrak, Nicholas E. (1989): Potential invasion of the Great Lakes by fish species associated with climatic warming. *Journal of Great Lakes Research* 15, 306-316. [FISH, GREAT LAKES]
- Manny; B.A.; Fahnenstiel, G.L.; Gardner, W.S. (1987): Acid Rain Stimulation of Lake Michigan.
- Marti, Edwin A.; Arrnstrong, David E. (1990): Polychlorinated biphenyls in Lake Michigan tributaries. *Journal of Great Lakes Research* 16, 396-405. [LAKE MICHIGAN, LOADINGS, PCB, POLLUTION, TOXICS, WATER COLUMN]
- Masalek, J. (1990): Evaluation of Pollutant Loads from urban Nonpoint Sources. *Water Science and Technology* 22, 23-30. [GREAT LAKES, LOADINGS, POLLUTION]
- Mansfield, P.J.; Jude, D.J.; Michaud, D.T.; Brazo, D.C.; Gulvas, J. (1983): Distribution and Abundance of Larval Burbot and Deepwater Sculpin in Lake Michigan. *Transactions of the American Fisheries Society* 112, Pages 162-172. [FISH, LAKE MICHIGAN]
- Mansfield, P.J.; Jude, D.J. (1986): Alewife (*Alosa pseudoharengus*) survival during the first growth season in southeastern Lake Michigan. Reprinted from *Canadian Journal of Fisheries and Aquatic Sciences*. Vol. 43, No. 7, pp 1318-1326. [FISH, LAKE MICHIGAN]
- Marshall, T.R. et al. (1987): Using the Lake Trout as an Indicator of Ecosystem Health: Application of the Dichotomous Key. (Great Lakes fishery Commission Report 49.) Ontario Ministry of Natural Resources, Canada. 41 pages. [ECOSYSTEM, FISH, GREAT LAKES, HABITAT, HEALTH, INDICATOR]
- Mass Balance Workshop Steering Committee (1991): Proceedings of the Mass Balance Workshop. Report to the Surveillance Subcommittee to the Great Lakes Water Quality Board. 91 pages. [GREAT LAKES, LAKE ONTARIO, MANAGEMENT, TOXICS]

May, R.M. (1981): Patterns in Multi-Species Communities. In: Theoretical Ecology: Principles and Applications. 2nd ed. (Ed: May, RM) Blackwell Scientific Publications, Oxford. [ECOSYSTEM]

McComas, Steve: LakeSmarts - The First Lake Maintenance Handbook. [FISH, SEDIMENTS, GREAT LAKES]

McComish, Thomas S.; Shroyer, Steven M. (1993): Characteristics of the Near - Shore Non-Salmonine Fish Community in Indiana Waters of Lake Michigan with Emphasis on Yellow Perch. Project Performance Report for 1992, Federal Aid Project F-18-R, Segment 5, Indiana Department of Natural Resources, Division of Fish and Wildlife. Pages.

Metropolitan Water Reclamation District of Greater Chicago (1982-86): Research and Development Report 93-2 of the Greater Chicago Water Reclamation District for Southwestern Lake Michigan. [WATER QUALITY, SEDIMENT QUALITY, BENTHIC INVERTEBRATES]

Metropolitan Water Reclamation District of Greater Chicago, Research and Development Department: Report on Lake Michigan Water Quality - Clean Water Quarterly: January Through March 1993. [LAKE MICHIGAN, WATER QUALITY]

Metropolitan Water Reclamation District of Greater Chicago, Research and Development Department: Report on Lake Michigan Water Quality - Clean Water Quarterly: April Through June 1993. [LAKE MICHIGAN, WATER QUALITY]

Metropolitan Water Reclamation District of Greater Chicago, Research and Development Department: Report on Lake Michigan Water Quality - Clean Water Quarterly: July Through September 1993. [LAKE MICHIGAN, WATER QUALITY]

Michigan Department of Natural Resources, Surface Water Quality Division (1986): Water Quality and Pollution Control in Michigan. Michigan 305(b) Report: Volume 9: Water Years 1984 and 1985. 132 Pages. [WATER QUALITY, LAKE MICHIGAN, GREAT LAKES, TOXICS, RIVERS, AQUATIC LIFE, FISH, HEALTH]

Michigan Department of Natural Resources, Office of the Great Lakes (1986): State of the Great Lakes: Annual Report for 1986. 40 Pages. [GREAT LAKES, LAKE MICHIGAN, SEDIMENTS, FISH, TOXICS]

Michigan Department of Natural Resources (1988): State of the Great Lakes 1978-1988. 36 pages. [GREAT LAKES, HUMAN USE, TRENDS]

Michigan Department of Natural Resources, Office of the Great Lakes: State of the Great Lakes 1988-1989. 48 Pages. [GREAT LAKES, LAKE MICHIGAN]

Michigan Department of Natural Resources, Office of the Great Lakes: Great Lakes Connections: State of the Great Lakes 1990-1991. 47 Pages. [GREAT LAKES, LAKE MICHIGAN, HUMAN HEALTH, FISH]

Michigan Department of Natural Resources (1990): Water Quality and Pollution Control in Michigan 1990 Report. Section 305(b) Report. 274 pages. [AVIAN SPECIES, EUTROPHICATION, FISH, HABITAT, HUMAN HEALTH, HUMAN USE, PLANKTON, SEDIMENT, TOXICS, WATER COLUMN, WETLAND]

Michigan Department of Natural Resources, Surface Water Quality Division (1992): Water Quality and Pollution Control in Michigan: 1992 Report. Michigan 305(b) Report: Volume 12. Lansing, MI. 351 Pages. [WATER QUALITY, LAKE MICHIGAN, GREAT LAKES, TOXICS, RIVERS, SEDIMENTS, AQUATIC LIFE, FISH, HEALTH]

Michigan Department of Natural Resources, Office of the Great Lakes (1993): State of the Great Lakes: 1993 Annual Report. 47 Pages. [GREAT LAKES, LAKE MICHIGAN, SEDIMENTS, FISH, TOXICS, HABITAT, CHLORINE]

Mills, Edward L.; Leach, Joseph H.; Carlton, James T.; Secor, Carol L. (1993): Exotic Species in the Great Lakes: A History of Biotic Crimes and Anthropogenic Introductions. Journal of Great Lakes Research 19, 1-54. [ECOSYSTEM, EXOTIC SPECIES, GREAT LAKES, TRENDS]

Mills, Edward L.; Leach, Joseph H.; Secor, Carol L.; Carlton, James T. (1993): What's Next? The Prediction and Management of Exotic Species in the Great Lakes (Report of the 1991 Workshop). Great Lakes Fishery Commission, Ann Arbor, MI. 22 pages. [EXOTIC SPECIES]

Milwaukee Metropolitan Sewerage District: Water Quality and Biological Data for Near Shore Lake Michigan - Information Provided to The ADVENT Group, Inc. on March 30, 1994. Jones Island Wastewater Treatment Plant. [LAKE MICHIGAN, WATER QUALITY]

Mlot, Christine (1989): Great Lakes fish and the greenhouse effect. BioScience 39(Mar), 145. [ATMOSPHERE, FISH, GREAT LAKES]

Moll, R. A.; Rossmann, R.; Bares, J.A.; Horvath, F.J. (1991): Historical Trends of Chlorides in the Great Lakes. In: Deicing Chemicals and the Environment (Ed: D'Itri, FM) Lewis Publishers, Chelsea, MI, 303-322. [GREAT LAKES, TOXICS]

Morehead, N.R.; Robbins, J.A.; Mudroch, A. (1991): Anthropogenic Alteration of Carbon, Nitrogen, and Biogenic Silica Deposition in Lake George (St Mary's River) Since 1820. In: 34 Annual Meeting of the International Association for Great Lakes Research, Buffalo, NY, June 2-6. [SEDIMENT, ST MARYS RIVER]

Mortimer, Clifford H.; Csanady, Gabriel T. (1975): Environmental Status of the Lake Michigan Region. Volume 2. Physical Limnology of Lake Michigan; Part 1. Physical Characteristics of Lake Michigan and its Responses to Applied Forces; Part 2. Diffusion and Dispersion. Argonne National Laboratory, Argonne, Illinois. [Hydrology, Wind Currents, Buoyancy, Seasonal Cycles]

Munawar, M.; Munawar, I.F. (1982): Toxicity Studies on Natural Phytoplankton Assemblages by Means of Fractionation Bioassays. Can. Fish. Aquat. Sci. Tech. Rep. No. 1152. [PLANKTON]

Munawar, M; Wong, PTS; Rhee, GY (1988): Effects of Contaminants on Algae: An Overview. In: Toxic Contamination in Large Lakes. Volume I: Chronic Effects of Toxic Contaminants in Large Lakes. Lewis Publishers, Chelsea, MI, 113-160. [GREAT LAKES, LAKE ONTARIO, LAKE SUPERIOR, PLANKTON, POLLUTION, TOXICS, WATER COLUMN]

Munawar, M.; Wiese, T. (1989): Is the "microbial loop" an early warning indicator of anthropogenic stress? *Hydrobiologia* 188-189, 163-174. [GREAT LAKES, INDICATOR]

Munawar, M. et al. (1989): Probing Ecosystem Health: a Multi-Disciplinary and Multi-Trophic Assay Strategy. *Hydrobiologia* 188-189, 93-116. [ECOSYSTEM, EUTROPHICATION, GREAT LAKES, HEALTH, NUTRIENTS, PLANKTON, SEDIMENT, TOXICS, WATER COLUMN]

Munkittrick, Kelly R., Dixon, D. George (1989): A Holistic Approach to Ecosystem Health Assessment Using Fish Population Characteristics. *Hydrobiologia* 188-189, 123-135. [ECOSYSTEM, FISH, HEALTH, INDICATOR, POPULATION]

Munkittrick, K.R.; Dixon, D.G. (1989): Use of White Sucker (*Catostomus commersoni*) Populations to Assess the Health of Aquatic Ecosystems Exposed to Low-Level Contaminant Stress. *Canadian Journal of Fisheries and Aquatic Sciences* 46, 1455-1462. [ECOSYSTEM, FISH, HABITAT, INDICATOR]

The Nature Conservancy (1994): The Conservation of Biological Diversity in the Great Lakes Ecosystem: Issues and Opportunities. The Nature Conservancy, Great Lakes Program. 118 Pages. [GREAT LAKES, AQUATIC LIFE, BIOLOGICAL DIVERSITY]

Nepszy, S.J.; Leach, J.H. (1973): First Records of the Chinese Mitten Crab, *Eriocheir sinensis* (Crustacea: Brachyrra) from North America. *Journal of the Fisheries Research Board of Canada* 30, 1909-1910. [EXOTIC SPECIES]

Nepszy, Stephen J. et al. (1988): Parasites of Fishes in the Canadian Waters of the Great Lakes. (Great Lakes fishery Commission Technical Report 51.). 112 pages. [FISH, GREAT LAKES]

NOAA, U.S. National Marine Pollution Program Office (1980): Report of Great Lakes Region Conference on Marine Pollution Problems Held at Traverse City Michigan, on June 9-11, 1980. Prepared for the Federal Coordinating Council for Science, Engineering, and Technology, Washington, DC. 73 Pages. (NTIS # PB82 106295) [TOXICS, GREAT LAKES, TOXICS, SEDIMENTS, HUMAN HEALTH]

Ohio EPA, Division of Water Quality, Planning and Assessment, Ecological Assessment Section: A Preliminary Assessment of Ohio's Lake Erie Estuarine Fish Communities. 27 Pages. [FISH, GREAT LAKES, RIVER]

Ohio EPA, Division of Water Quality, Planning and Assessment, Ecological Assessment Section (1990): Ohio Water Resource Inventory, Executive Summary and Volume 1. 174 Pages. [FISH, GREAT LAKES, RIVER, STREAMS, BIOASSAY, AQUATIC LIFE, WATER QUALITY STANDARDS, HEALTH, HABITAT, TOXICS]

Ohio EPA, Division of Water Quality, Planning and Assessment, Ecological Assessment Section (1990): Compendium of Biological Results From Ohio Rivers, Streams, and Lakes: 1989 Edition. 176 Pages. [FISH, GREAT LAKES, RIVER, STREAMS, HABITAT]

Perrone, Michael Jr.; Schneeberger, Philip J.; Jude, David J. (1983): Distribution of Larval Yellow Perch (*Perca Flavescens*) in Nearshore Waters of Southeastern Lake Michigan. J. Great Lakes Res. 9(4), Pages 517-522. [FISH, LAKE MICHIGAN]

Phytoplankton Growth. Journal of Great Lakes Research 13, 218-223. [ATMOSPHERE, ECOSYSTEM, LAKE MICHIGAN, POLLUTION, WATER COLUMN]

Polls, I.; Sedita, S.J.; Zenz, D.R.; Lue-Hing, C. (1993): Research and Development Department - Report No. 93-2: A Comparison of the Water and Sediment Quality and Benthic Invertebrates in the Grand Calumet River, the Indiana Harbor Canal, Indiana Harbor, Southwestern Lake Michigan, and the Calumet River During 1982 and 1986. Metropolitan Water Reclamation District of Greater Chicago. [WATER QUALITY, SEDIMENT, BENTHIC]

Pontaseh, K.W.; Smith, E.P.; Cairns, Jr., J. (1989): Diversity indices, community comparison indices and canonical discriminant analysis: Interpreting the results of multispecies toxicity tests. Water Resources 23, 1229-1238. [INDICATOR, TOXICS]

Predation and Weather on Long-Term Water Quality Trends in Lake Michigan. Canadian Journal of Fisheries and Aquatic Sciences 43, 435-443. [FISH, LAKE MICHIGAN, TRENDS]

Pringle, C.M.; White, D.S.; Rice, C.P.; Tuchman, M.L. (1981): The Biological Effects of Chloride and Sulfite with Special Emphasis on the Laurentian Great Lakes. Publ. No. 20. Great Lakes Research Div. University of Michigan, Ann Arbor. 51 pages. [GREAT LAKES, POLLUTION]

Project Management Team, Functional Group 2 (1989): Living With the Lakes: Challenges and Opportunities. Annex B: Environmental Features, Processes and Impacts - An Ecosystem Perspective on the Great Lakes-St Lawrence River System, Burlington, Ontario and Chicago, Illinois. 169 pages. [ECOSYSTEM, GREAT LAKES, MANAGEMENT]

Project Management Team, Functional Group 5 (1989): Living With the Lakes: Challenges and Opportunities. Annex D: The Great Lakes Ecosystem Perspective - Implications for Water Levels Management, Burlington, Ontario and Chicago, Illinois. 79 pages. [ECOSYSTEM, GREAT LAKES, MANAGEMENT]

Public Health Service, Washington, D.C. (March 2, 1965): Pollution Of The Interstate Waters Of The Grand Calumet River, Little Calumet River, Calumet River, Wolf Lake, Lake Michigan and Their Tributaries. Proceedings Of Conference held at Chicago, Illinois on March 2-9, 1965. Volume I. (PB-230 544)

Public Health Service, Washington, D.C. (March 3, 1965): Pollution Of The Interstate Waters Of The Grand Calumet River, Little Calumet River, Calumet River, Wolf Lake, Lake Michigan And Their Tributaries. Proceedings of Conference held at Chicago, Illinois on March 2-9, 1965. Volume 2. (PB-230 545)

Public Health Service, Washington, D.C. (March 4, 1965): Pollution Of The Interstate Waters Of The Grand Calumet River, Little Calumet River, Calumet River, Wolf Lake, Lake Michigan And Their Tributaries. Proceedings of Conference held at Chicago, Illinois on March 2-9, 1965. Volume 3. (PB-230 546)

Public Health Service, Washington, D.C. (March 5, 1965): Pollution Of The Interstate Waters Of The Grand Calumet River, Little Calumet River, Calumet River, Wolf Lake, Lake Michigan And Their Tributaries. Proceedings of Conference held at Chicago, Illinois on March 2-9, 1965. Volume 4. (PB-230 547)

Quinn, John P.; Janssen, John. (1989): Crayfish Competition in Southwestern Lake Michigan: A Predator Mediated Bottleneck. Journal of Freshwater Ecology, Volume 5, Number 1. [PLANKTON, LAKE MICHIGAN, BENTHIC]

Rabe, Barry G.; Zimmerman, Janet B. (1992): Cross-Media Environmental Integration in the Great Lakes Basin. Environ Law 22, 253-289. [GREAT LAKES, MANAGEMENT, POLLUTION]

Rakoczy, G.P.; Rogers, R.D. (1987): Sportfishing Catch and Effort from the Michigan Waters of Lakes Michigan, Huron, and Erie, and their Important Tributary Streams, April 1, 1986-March 31, 1987. Michigan Department of Natural Resources, Fisheries Division Fisheries Technical Report Number 87-6A. [FISH, HUMAN USE, LAKE ERIE, LAKE HURON, LAKE MICHIGAN]

Rathke, David E.; and McRae, Gil (1989): 1987 Report on Great Lakes Water Quality, Appendix B Great Lakes Surveillance, Volume I.

Rathke, David E.; and McRae, Gil (1989): 1987 Report on Great Lakes Water Quality, Appendix B Great Lakes Surveillance, Volume II.

Rathke, David E.; and McRae, Gil (1989): 1987 Report on Great Lakes Water Quality, Appendix B Great Lakes Surveillance, Volume III.

Reed, P., Evans, R.L. (1981): Acute Toxicity of Chlorides, Sulfates, and Total Dissolved Solids to Some Fishes in Illinois. 50 p. Illinois State Water Survey, Office of Publications Services, Illinois Department of Energy and Natural Resources.

Richardson, W.L.; Paul, J.F. (1988): Historical Synopsis of Great Lakes Water Quality Research and Management and Future Directions. In: Protection of River Basins, Lakes, and Estuaries, Section 2, Protection and Management of Water Quality in Lakes and Estuaries. 83-97. (EPA/600/D-89/230, ERLN-1114) [DDT, EUTROPHICATION, GREAT LAKES, LAKE MICHIGAN, PCB, PLANKTON, POLLUTION, SEDIMENT, TOXICS, TRENDS, WATER COLUMN]

Robbins, J.A.; Morehead, N.R.; Mudroch, A. (1992): History of Contamination of Lake George and its Impact on Biogenic Silica Production from 1750-1986. In: Annual Meeting of the American Society of Limnology and Oceanography, Santa Fe, MN, February 9-14. [TOXICS]

Robertson, A. (1984): The Present Status of Research on the Zooplankton and Zoobenthos of the Great Lakes. *Journal of Great Lakes Research* 10, 156-163. [BENTHOS, GREAT LAKES, PLANKTON]

Rockwell, D.C. et al. (1980) Lake Michigan Intensive Study, 1976-1977. USEPA: December, 186 pp. (EPA-905/4-80-003-A) [LAKE STUDIES]

Rockwell, D.C.; Salisbury, D.K.; Lesht, B.M. (1989): Water Quality in the Middle Great Lakes: Results of the 1985 U.S. EPA (Environmental Protection Agency) Survey of Lake Erie, Huron, and Michigan: Final Report - Environmental Protection Agency, Chicago, IL Great Lakes National Program Office. 270 pages. (EPA/905/6-89/001, GLNPO-4) [EUTROPHICATION, GREAT LAKES, LAKE ERIE, LAKE HURON, LAKE MICHIGAN, NITRATE, PHOSPHORUS, POLLUTION, WATER COLUMN]

Rodgers, P.W.; Salisbury, D.K. (1981): Water Quality Modeling of Lake Michigan and Consideration of the Anomalous Ice Cover of 1976-1977. *Journal of Great Lakes Research* 7, 467-480. [LAKE MICHIGAN, WATER COLUMN]

Rodgers, Paul W.; Dilks, David W.; Samson, P. (1991). An integrated Model of Atmospheric and Aquatic Chemical Fate Useful for Guiding Regulatory Decisions: A Proposal. In: Long Range Transport of Pesticides. Kurtz, David A. (ed). Lewis Publishers, Chelsea, MI. [ATMOSPHERE, TOXICS]

Romano, R.R.; McIntosh, A.W.; Kessler, W.V. et.al. (1977): Trace Metal Discharges of the Grand Calumet River. *J. Great Lakes Res.* October 1977. *Internat. Assoc. Great Lakes Res.* 3(1-2):144-147. [LAKE MICHIGAN]

Rossmann, R. (1980): Soluble Element Concentrations and Complexations in Southeastern Lake Michigan: *Journal of Great Lakes Research* 6, 47-53. [LAKE MICHIGAN]

Rossmann, R.: Trace Metal Concentrations in the Offshore Waters of Lakes Erie and Michigan. Special Report No. 108. Great Lakes Research Division: November 1984, 190 pp. (NTIS PB-85-199396-AS).

Rossmann, Ronald; Chang, William Y. B.; Bowers, James A.; Feldt, Laurie; Barres, James: Southeastern Nearshore Lake Michigan: Impact of the Donald C. Cook Nuclear Plant. University of Michigan, Great Lakes Research Division, Publication 22.

Rossmann, Ronald; Barres, James (1988): Trace Element Concentration in Near-Surface Waters of the Great Lakes and Methods of Collection, Storage, and Analysis. *Journal of Great Lakes Research* 14, 188-204. [GREAT LAKES, LAKE ERIE, LAKE HURON, LAKE MICHIGAN, LAKE ONTARIO, LAKE SUPERIOR, LEAD, MERCURY, TOXICS, WATER COLUMN]

Rowan, David J.; Rasmussen, Joseph B. (1992): Why don't Great Lakes fish reflect environmental concentrations of organic contaminants? - An analysis of between-lake variability in the ecological partitioning of PCBs and DDT. *Journal of Great Lakes Research* 18, 724-741. [DDT, ECOSYSTEM; FISH, GREAT LAKES, PCB, TOXICS]

Ryder, R.A.; Edwards, C.J. (Eds.) (1985): A Conceptual Approach for the Application of Biological Indicators of Ecosystem Quality in the Great Lakes Basin Report to the Great Lakes Science Advisory Board of the International Joint Commission., Windsor, ON. 169 pages. [ECOSYSTEM, GREAT LAKES, INDICATOR]

Ryder, Richard A.; Kerr, Steve R. (1990): Aquatic harmonic communities: surrogates of ecosystem integrity. In: An Ecosystem Approach to the Integrity of the Great Lakes in Turbulent Times. Proceedings of a 1988 Workshop Supported by the Great Lakes Fishery Commission and the Science Advisory Board of the International Joint Commission. Great Lakes Fishery Commission Seal Publication 90-4. (Eds:Edwards,CJ; Regier, Henry,A) Great Lakes Fishery Commission, Ann Arbor, MI, 239-255. [ECOSYSTEM; GREAT LAKES, INDICATOR, WILDLIFE]

Ryder, R.A. (1990): Ecosystem health, a human perception: definition, detection, and the dichotomous key. Journal of Great Lakes Research 16,619-624. [ECOSYSTEM, GREAT LAKES, HEALTH, INDICATOR]

Samples, K.; Bishop, R. (1981): The Lake Michigan Angler: A Wisconsin Profile. University of Wisconsin Sea Grant Institute Pub. WIS-SG-81-423., Madison, WI. 59 pages. [FISH, HUMAN USE, LAKE MICHIGAN]

Scavia, Donald et al. (1988): Dynamics of Lake Michigan Plankton: a Model Evaluation of Nutrient Loading, Competition, and Predation. Canadian Journal of Fisheries and Aquatic Sciences 45, 165-177. [EUTROPHICATION, LAKE MICHIGAN, PLANKTON, TRENDS, WATER COLUMN]

Scavin, D.; Fahnenstiel, G.; Evans, M.; Jude, D.; Lehman, J. (1986): Influence of Salmonine Predation and Weather on Long-Term Water Quality Trends in Lake Michigan. Canadian Journal of Fisheries and Aquatic Sciences 43, 435-443. [FISH, LAKE MICHIGAN, TRENDS]

Schelske, Claire, L.; Stoermer, Eugene F. (1971): Eutrophication, Silica Depletion, and Predicted Changes in Algal Quality in Lake Michigan. [EUTROPHICATION, SILICA, PHYTOPLANKTON]

Schelske, C.L.; Feldt, L.E.; Simmons, M.S. (1980): Phytoplankton and Physical Chemical Conditions in Selected Rivers and the Coastal Zone of Lake Michigan, 1972 Publ. No 19. Great Lakes Res. Div., Univ. of Michigan, Ann Arbor. 162 pages. [LAKE MICHIGAN, PLANKTON]

Schelske, C.L.; Stoermer, E.F.; Gannon, J.E. and Simmons, M.S. (1976): Biological, chemical and physical relationships in the Straits of Mackinaw. University of Michigan, Great Lakes Research Div. Spec. Rept. 60, 267 pages. -

Schelske, C.L.; Feldt, L.E.; Simmons, M.S.; and Stoermer, E.F. (1974): Storm induced relationships among chemical conditions and phytoplankton in Saginaw Bay and Western Lake Huron. Proc. 17th Conf. Great Lakes Res., pp. 78-91. International Association Great Lakes Research.

Schelske, C.L.; Moll, R.A.; Berry, T.; Stoermer, E.F. (1985): Limnological Characteristics of Northern Lake Michigan, 1976. Spec. Rep. No. 95. Great Lakes Research Division, Univ. of Michigan, Ann Arbor. 245 pages. [LAKE MICHIGAN]

Schelske, C.L.; Stoermer, E.F.; and Feldt, L.E. (1971): Nutrients, phytoplankton productivity, and species composition as influenced by upwelling in Lake Michigan. Proc. 14th Conference. Great Lakes Research, International Association Great Lakes Research. pp.102-113.

Schelske, C.L.; Feldt, L.E.; Santiago, M.A. and Stoermer, E.F.; (1972): Nutrient enrichment and its effect on phytoplankton and species composition in Lake Superior. Proc. 15th Conf. Great Lakes Res., Int. Assoc. Great Lakes Res., pp. 149-165.

Schelske, C.L.; and Stoermer, E.F.; (1972): Phosphorus, silica and eutrophication in Lake Michigan. In G.E. Likens (ed.). Nutrients and Eutrophication: The Limiting-nutrient Controversy. Special Symposium. Vol. 1, ASLO, pp. 157-171.

Schelske, C.L.; Stoermer, E.F.; Fahnenstiel, G.L.; Haibach, M. (1986): Phosphorus Enrichment Silica Utilization, and Biogeochemical Silica Depletion in the Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences 43, 407-415. (EPA/600/J-86/522 Grants EPA-R406294, NSF-OCE82-16588) [EUTROPHICATION, GREAT LAKES, LAKE ERIE, LAKE ONTARIO, PHOSPHORUS, WATER COLUMN]

Schelske, Claire L. (1991): Historical Nutrient Enrichment of Lake Ontario: Paleolimnological Evidence Canadian Journal of Fisheries and Aquatic Sciences 48, 1529-1538. [EUTROPHICATION, LAKE ONTARIO, LOADINGS, NUTRIENTS, PHOSPHORUS, SEDIMENT, TRENDS, WATER COLUMN]

Schierow, L.J.; Chesters, G. (1988): Evaluation of the Great Lakes Near Shore Index. Water Research 22, 269-277. [EUTROPHICATION, GREAT LAKES, MANAGEMENT, POLLUTION, WATER COLUMN]

Schindler, D.W. (1987): Detecting ecosystem responses to anthropogenic stress. Can. J. Fish: Aquat. Sci. 44(Suppl 1), 6-25.[ECOSYSTEM]

Schneider, J.C.; Leach, J.H. (1979): Walleye Stocks in the Great Lakes 1800-1975: Fluctuations and Possible Causes. Technical Report No. 31. Great Lakes Fishery Commission, Ann Arbor, MI.[FISH] Research 22, 269-277. [EUTROPHICATION, GREAT LAKES, MANAGEMENT, POLLUTION, WATER COLUMN]

Schneider, J.C. et al. (1991): Walleye Rehabilitation in Lake Michigan, 1969-1989. In: Great Lakes Fishery Commission Special Publication 91-1. 23-61.[FISH, LAKE MICHIGAN, MANAGEMENT, POPULATION, RESTORATION, TRENDS, WATER COLUMN]

Science Applications International Corporation (1993): Revised Lake Michigan Lakewide Management Plan for Toxic Pollutants (Draft). 225 Pages. U.S. EPA Region V, Chicago IL. (EPA Contract # 68-C8-0066, W.A. No. C-4-98(O). [LAKE MICHIGAN])

Sediment Concentrations from Multiple Day Landsat and AVHRR data. *Journal of Remote Sensing of the Environment* 24, 364-373. [SEDIMENT]

Serafin, Rafal (1990): Rehabilitating Great Lakes Integrity in Times of Surprise. In: *An Ecosystem approach to the Integrity of the Great Lakes in Turbulent Times* (Great Lakes Fishery Comm 90-4). 45-67.[ECOSYSTEM, GREAT LAKES, RESTORATION]

Sheath, R.G. (1987). Invasions into the Laurentian Great Lakes by Marine Algae. *Arch. Hydrobiol. Beih. rgebn. Limnol.* 25, 165-186.[EXOTIC SPECIES, GREAT LAKES]

Simmers, J.W.; Lee, C.R.; Brandon, D.L.; Tatem, H.E.; Skogerboe, J.G. (1991): Information Summary, Area of Concern: Grand Calumet River, Indiana. 203 Pages. Miscellaneous Paper EL-91-10. USAE Waterways Experiment Station, Environmental Laboratory, Vicksburg, MS. [TOXICS, SEDIMENTS, GREAT LAKES, WATER QUALITY CRITERIA]

Smith, S.H. (1972): Factors in Ecologic Succession in Oligotrophic Fish Communities of the Laurentian Great Lakes. *Journal of the Fisheries Research Board of Canada* 29, 717-730.[FISH; GREAT LAKES]

Sonzogni, W .L.; Rodgers, P.W.; Richardson, W. L; Monteith, T. J. (1982). Chloride Pollution of the Great Lakes. *Journal of the Water Pollution Control Federation.* 55(5), 513-521. [GREAT LAKES, POLLUTION]

Sonzogni, W.C.; Robertson, A.; Beeton, A.M. (1983): Great Lakes Management: Ecological Factors. *Environmental Management* 7, 531-542.[GREAT LAKES, MANAGEMENT]

Sonzogni, William C; Heidtke, Thomas M (1986): Water Quality Management for the Great Lakes. *Journal of Water Resources Planning and Management* 1 12(Jan), 48-63. [GREAT LAKES, MANAGEMENT, POLLUTION, WATER COLUMN]

Sonzogni, W.C.; Canale, R.P.; Larn, DCL.; Lick, W.; Mackay, D. (1987): Large Lake Models'- Uses, Abuses, and Future. *Journal of Great Lakes Research* 13, 387-396. (Environmental Research, Lab - Duluth, Grosse Ile, MI. Large Lakes Research Station). [EUTROPHICATION, GREAT LAKES, POLLUTION, TOXICS, WATER COLUMN]

Sprules, W. Gary; Riessen, Howard P.; Jin, Edward H. (1990): Dynamics of the Bythotrephes invasion of the St Lawrence Great Lakes. *Journal of Great Lakes Research* 16, 346-351. [EXOTIC SPECIES, GREAT LAKES]

Stanley, J.G.; Peoples, R.A. Jr; Mcann, J.A. (1991): U.S. Federal Policies, Legislation, and Responsibilities Related to Importation of Exotic Fishes and Other Aquatic Organisms. *Canadian Journal of Fisheries and Aquatic Sciences* 45 (Suppl 1), 162-166.[EXOTIC SPECIES]

Stearns, Forest; Hole, Francis, D.; Klopatek, Jeffery (1974): Environmental Status of the Lake Michigan Region. Volume 9. *Soils of the Lake Michigan Drainage Basin-An Overview.* Argonne National Laboratory, Argonne, Illinois.

- Stearns, Forest; Hole, Lindsley, Diane (1977): Environmental Status of the Lake Michigan Region. Volume 11. Natural Areas of the Lake Michigan Drainage Basin and Endangered or Threatened Plant and Animal Species. Argonne National Laboratory, Argonne, Illinois
- Steedman, Robert J.; Regier, Henry A. (1990): Ecological bases for an understanding of ecosystem integrity in the Great Lakes basin. In: An Ecosystem Approach to the Integrity of the Great Lakes in Turbulent Times. Proceedings of a 1988 Workshop Supported by the Great Lakes Fishery Commission and the Science Advisory Board of the International Joint Commission. Great Lakes Fishery Commission Special Publication 90-4. (Eds: Edwards, C.J.; Regier, Henry, A.) Great Lakes Fishery Commission, Ann Arbor, MI, 257-270. [ECOSYSTEM, GREAT LAKES, HUMAN USE]
- Stemberger, R.S., and M. S. Evans. (1984): Rotifer Composition and Abundance in Lake Ontario During IFYGL. National Environmental Research Center.
- Stemberger, R.S., and M. S. Evans. (1984): Rotifer seasonal succession and copepod predation in Lake Michigan. J. Great Lakes Res. 10(4): 417-428.
- Stoermer, E.F., and Ladewski, T.B. (1975): Apparent optimal temperatures for the occurrence of some common phytoplankton species in southern Lake Michigan. Great Lakes Res. Div., Univ. Michigan Publ. 18, 49 p.
- Stoermer, Eugene F.; and Yang, J.J. (1970): Distribution and relative abundance of dominant plankton diatoms. Univ. Michigan, Great Lakes Res. Div., Pub. No. 16. 64 p.
- Stoermer, Eugene F.; Conley, Daniel J.; Schelske, Claire L. (1983): Early eutrophication in the lower Great Lakes: new evidence from biogenic silica in sediments. Science 222(21 Oct), 320-322. [EUTROPHICATION, GREAT LAKES, SEDIMENT, WATER COLUMN]
- Stoermer, Eugene F.; Stevenson, R.J. (1979): Green Bay phytoplankton, composition, abundance and distribution. EPA-905/3-79-002.
- Stoermer, E.F., Tuckman, M.L. (1979): Phytoplankton Assemblages of the Near-Shore Zone of Southern Lake Michigan. USEPA, November, 99 pp. (EPA 905/2-87-002). [AQUATIC LIFE: FISH STUDIES AND OTHER BIOTA]
- Stoermer, E.F. (1978): Phytoplankton Assemblages as indicators of water quality in the Laurentian Great Lakes. Trans. Amer. Micros. Soc. 97 (1): 2-16.
- Stoermer, E.F., et al (1975): Phytoplankton Composition and Abundance in Lake Ontario During IFYGL. National Environmental Research Center.
- Stoermer, E.F., and Theriot, E. (1985): Phytoplankton distribution in Saginaw Bay. J. Great Lakes Res. 11 (2): 132-142.
- Stoermer, E.F., and Kopczynska, E. (1967a): Phytoplankton populations in the extreme southern basin of Lake Michigan, 1962-1963. Proc. 10th Conf. Great Lakes Res., Int. Assoc. Great Lakes Res. pp. 88-106.

- Stoermer, E.F., and Kopczynska, E. (1967b): Phytoplankton populations in the extreme southern basin of Lake Michigan, 1962-1963, pp. 19-40. In J.C. Ayers and D.C. Chandler. Studies on the environment and eutrophication of Lake Michigan. Univ. Michigan, Great Lakes Res. Div., Spec. Rep. No. 30.
- Stoermer, E.F., Schelske, C.L., Santiago, M.A. and Feldt, L.E. (1972): Spring phytoplankton abundance and productivity in Grand Traverse Bay, Lake Michigan, 1970. 15th Conf. Great Lakes Res. 1972: 181-191. Intern. Assoc. Great Lakes Res.
- Strachan, WM.J. (1988): Test systems and exposure in the aquatic Environment Ambio 17, 394-397.[ECOSYSTEM, GREAT LAKES, RISK ASSESSMENT]
- Sullivan, R.A.C., Baise, M., Sonzogni, W.C. (1981): Environmental Quality Maps for the U.S. Great Lakes Basin. USEPA, June, 98 pp. (EPA 905/4-79-028. [GREAT LAKES BASIN]
- Task Force on Chemical Loadings of the Toxic Substances Committee (1988): Report on Modeling the Loading-Concentration Relationship for Critical Pollutants in the Great Lakes. Report to the Great Lakes Water Quality Board., Windsor, Ontario. 275 pages. [GREAT LAKES, LOADINGS, MANAGEMENT, PCB, POLLUTION, TOXICS, WATER COLUMN]
- Thomann, Robert V.; Mueller, John A. (1987). Principles of Surface Water Quality Modeling and Control. Harper and Row, Publishers, New York. 644 pages. [GREAT LAKES]
- Tin, Heang, T.; and Jude, David, J. (1983): Distribution and Growth of Larval Rainbow Smelt in Eastern Lake Michigan, 1978-1981.
- Tissue, Thomas; Edgington, David N.; Seils, Charles A. (1988): Sulfate Reduction in Sediment Interstitial Fluids in Lakes Michigan and Erie. Journal of Great Lakes Research 14, 14-22. [EUTROPHICATION, LAKE ERIE, LAKE MICHIGAN, NUTRIENTS, TRENDS]
- Torrey, Marguerite S. (1976): Environmental Status of the Lake Michigan. Chemistry of Lake Michigan Argonne National Laboratory, Argonne, Illinois. ANL/ES-40 Volume 3. [Conventional Pollutants, Nutrients, Data Review, Criteria]
- The University of Michigan, Great Lakes Research Division (1979): Phytoplankton Assemblages of the Nearshore Zone of Southern Lake Michigan. 88 Pages. Prepared for U.S. EPA Great Lakes National Program Office, Chicago, IL. (Grant No.- R005337-001, NTIS # PB81-106387). [PLANKTON, LAKE MICHIGAN]
- The University of Michigan, Great Lakes and Marine Waters Center. (1984): Field Distribution and Entrainment of Fish Larvae and Eggs at the Donald C. Cook Nuclear Power Plant, Southeastern Lake Michigan, 1973-1979. Special Report No. 105 of the Great Lakes Research Division. 320 Pages. [FISH, LAKE MICHIGAN]
- The University of Michigan, Great Lakes and Marine Waters Center. (1985): Zooplankton Studies at the Donald C. Cook Nuclear Plant: 1979-1982 Investigations, Including Preoperational (1971-1974) and Operational (1975-1982) Comparisons. Special Report No. 111 of the Great Lakes Research Division. 236 Pages. [PLANKTON, LAKE MICHIGAN]

U.S. Army Corps. of Engineers (1986): Indiana Harbor Combined Disposal Facility and Maintenance Dredging, Lake Co., IN Draft EIS, February. [SEDIMENT, WASTE DISPOSAL]

U.S. Army Corps of Engineers, et. al., A Joint Federal/State 5-Year Strategy (1992-1997), Protecting the Great Lakes - Our Environmental Goals and How We Plan to Achieve Them. April 1992 DRAFT.

U.S. Army Corps of Engineers, U.S. Environmental Protection Agency, National Oceanographic and Atmospheric Administration, U.S. Coast Guard, U.S. Department of Agriculture,

U.S. Fish and Wildlife Service, - IL, IN, MI, MN, NY, OH, PA, WI, and Chippewa/Ottawa Treaty Fishery Management Authority. A Joint Federal/State 5-Year Strategy (1992-1997): Protecting The Great Lakes - Our Environmental Goals and How We Plan to Achieve Them (April 1992 DRAFT).

U.S. Department of Health, Education, and Welfare, Technical Committee on Water Quality (1970): Water Quality in the Calumet Area. Conference on Pollution of Lower Lake Michigan, Calumet River, Grand Calumet River, Little Calumet River, and Wolf Lake, Illinois and Indiana. 131 Pages. (NTIS # PB245377). [LAKE MICHIGAN, RIVERS, INDUSTRIAL DISCHARGE, WATER QUALITY]

U.S. EPA, Office of Water Regulations and Standards (1983): Technical Guidance Manual for Performing Waste Load Allocations. Book II - Streams and Rivers: Chapter 1 - BOD/DO. (EPA D-337)(Contract No. 68-01-5918).

U.S. EPA, Office of Water Regulations and Standards (1983): Technical Guidance Manual for Performing Waste Load Allocations. Book IV - Lakes and Impoundments: Chapter 2 Nutrient/Eutrophication Impacts. 179 Pages. (EPA 440/4-84-019)

U.S. EPA, Office of Water (1983): Technical Support Manual: Waterbody Surveys and Assessments for Conducting Use Attainability Analysis. 220 Pages. Washington D.C. [CHEMICAL, PHYSICAL, BIOLOGICAL, HABITAT, RIVERS, STREAMS]

U.S. EPA, Office of Water (1983): Technical Support Manual: Waterbody Surveys and Assessments for Conducting Use Attainability Analysis - Volume II: Estuarine Systems. 185 Pages. Washington D.C. [CHEMICAL, PHYSICAL, FISH, PLANKTON, BENTHOS, AQUATIC LIFE]

U.S. EPA, Office of Water (1984): Technical Support Manual: Waterbody Surveys and Assessments for Conducting Use Attainability Analysis - Volume III: Lake Systems. 196 Pages. Washington D.C. [CHEMICAL, PHYSICAL, FISH, PLANKTON, BENTHOS, AQUATIC LIFE]

U.S. EPA, Office of Water Regulations and Standards (1984): Technical Guidance Manual for Performing Waste Load Allocations. Book II - Streams and Rivers: Chapter 3 - Toxic Substances. 404 Pages. (EPA 440/4-84-022)

- U.S. EPA, Office of Public Affairs (1985): The Great Lakes. EPA Journal, Vol. 11 No.2, March 1985. 32 Pages. [GREAT LAKES, TOXICS, SEDIMENTS]
- U.S. EPA, Office of Regulation and Standards Monitoring and Data Support Division (1986): Technical Guidance Manual for Performing Waste Load Allocations. Book IV - Lakes, Reservoirs and Impoundments: Chapter 3 - Toxic Substances Impact. 80 Pages. (EPA 440/4-87-002), Washington D.C.
- U.S. EPA, Office of Water Regulations and Standards (1986): Technical Guidance Manual for Performing Waste Load Allocations. Book VI - Design Conditions: Chapter 1 - Stream Design Flow for Steady State Modelling. 59 Pages. (EPA 440/4-86 014)
- U.S. EPA, Chicago, IL Great Lakes National Program Office (1988): Five Year Program Strategy for the Great Lakes National Program Office, FY1989-1993. 94 pages. (EPA/905/9-89/001, GLNPO 89/1) [EUTROPHICATION, FISH, GREAT LAKES, MANAGEMENT, PROGRAM, SEDIMENT, TOXICS, WATER COLUMN]
- U.S. EPA, Great Lakes National Program Office (1988): Phytoplankton and Zooplankton in Lakes Erie, Huron, and Michigan: 1984. 275 Pages. (EPA 905/3-88-001, GLNPO 03-88). Chicago, IL. [PLANKTON, GREAT LAKES, LAKE MICHIGAN]
- U.S. EPA, Office of Water Regulations and Standards, Criteria and Standards Division (1988): Ambient Water Quality Criteria for Chloride.
- U.S. EPA, Chicago, IL Great Lakes National Program Office (1989): U. S. Progress in Implementing the Great Lakes Water Quality Agreement: Annex Reports to the International Joint Commission, 1988. Final rept 89 pages. (EPA/905/9-89/006, GLNPO-05/89) [ATMOSPHERE, EUTROPHICATION, GREAT LAKES, GREEN BAY, GROUNDWATER, PHOSPHORUS, POLLUTION, SEDIMENT, TOXICS, WATER COLUMN]
- U.S. EPA, Great Lakes National Program Office (1989): Phytoplankton and Zooplankton in Lakes Erie, Huron, and Michigan: 1985. 252 Pages. (EPA 905/3-90-003, GLNPO 01-91). Chicago, IL. [PLANKTON, GREAT LAKES, LAKE MICHIGAN]
- U.S. EPA, Office of Water (1990): Technical Guidance Manual for Performing Waste Load Allocations. Book III - Estuaries: Part 1 - Estuaries and Waste Load Allocation Models. 65 Pages. (EPA 823-R-92-002), Washington D.C.
- U.S. EPA, Office of Water (1990): Technical Guidance Manual for Performing Waste Load Allocations. Book III - Estuaries: Part 2 - Application of Estuarine Waste Load Allocation Models. 137 Pages. (EPA 823-R-92-003), Washington D.C.
- U.S. EPA (1991): Development of Index of Biotic Integrity Expectations for the Ecoregions of Indiana, 1. Central Corn Belt Plain.
- U.S. EPA, Region V (1991): Development of Index of Biotic Integrity Expectations for the Ecoregions of Indiana - I. Central Corn Belt Plan. 93 Pages. (EPA 905/9-91/025). [WATER QUALITY, FISH]

- U.S. EPA, Environmental Research Laboratory, Office of Research and Development (1992): MICHTOX: A Mass Balance and Bioaccumulation Model for Toxic Chemicals in Lake Michigan. First Draft, July 4, 1992. [BIOACCUMULATION, LAKE MICHIGAN]
- U.S. EPA (1992): Exposure and Effects of Airborne Contamination for the Great Waters Program Report. 201 pages. [ATMOSPHERE, TOXICS]
- U.S. EPA, Office of Water (1992): Technical Guidance Manual for Performing Waste Load Allocations. Book III - Estuaries: Part 3 - Use of Mixing Zone Models in Estuarine Waste Load Allocations. 41 Pages. (EPA 823-R-92-004)
- U.S. EPA, Office of Water (1992): Technical Guidance Manual for Performing Waste Load Allocations. Book III - Estuaries: Part 4 - Critical Review of Coastal Embayment and Estuarine Waste Load Allocation Modeling. 69 Pages. (EPA 823-R-92-005)
- U.S. EPA. (1993): Revised Draft Lake Michigan Lakewide Management Plan for Toxic Pollutants. [LAKE MICHIGAN, MANAGEMENT]
- U.S. EPA, Great Lakes National Program Office (1993): Assessment and Remediation of Contaminated Sediments (ARCS) Program - Biological and Chemical Assessment of Contaminated Great Lakes Sediment. 351 Pages. (EPA-905-R93-006). Chicago, IL[SEDIMENTS, GREAT LAKES, BENTHOS]
- U.S. EPA, Office of Water, Water Quality Standards Branch (1993): Water Quality Standards Handbook, Second Edition. 646 Pages. (EPA 823-B-93-002). Washington D.C. [WATER QUALITY, STANDARDS, USE ATTAINABILITY, WATER QUALITY CRITERIA]
- U.S. EPA, Office of Water, Office of Science and Technology (1993): Water Quality Standards Academy: Basic Course: Participant Manual. 530 Pages. [WATER QUALITY, USE ATTAINABILITY, RISK ASSESSMENT, HEALTH, SEDIMENTS, AQUATIC LIFE]
- Vallentyne, John R., Beeton, Alfred M. (1988): The 'Ecosystem' Approach to Managing Human Uses and Abuses of Natural Resources in the Great Lakes Basin. Environmental Conservation 15, 58-62.[ECOSYSTEM, GREAT LAKES, MANAGEMENT, WATER COLUMN]
- Virtual Elimination Task Force of the International Joint Commission (1991): Persistent Toxic Substances: Virtually Eliminating Inputs to the Great Lakes. 42 pages. [AVIAN SPECIES, DDT, DEFORMITIES, DIOXIN, FISH, GREAT LAKES, HEALTH HUMAN HEALTH, LEAD, MAMMALIAN SPECIES, MERCURY, ORGANOCHLORINE, PCB, PESTICIDE, POLLUTION, POPULATION, PROGRAM; TOXICS, WILDLIFE]
- Voldner, E.C.; Alvo, M. (1989): Estimation of Sulphur and Nitrogen Wet Deposition to the Great Lakes. Environmental Science and Technology 23, 1223-1232. [ATMOSPHERE, GREAT LAKES]

- Voldner, E.C.; Smith, L. (1991): Production, Usage, and Atmospheric Emissions of 14 Priority Toxic Chemicals. Appendix 2 of the Proceedings of the Workshop on Great Lakes Atmospheric Deposition, October 29-31, 1986. Report to the Great Lakes Water Quality, Great Lakes Science Advisory and the International Air Quality Advisory Boards of the International Joint Commission., Windsor, Ontario. 100 pages. [ATMOSPHERE, GREAT LAKES, LOADINGS, TOXICS]
- Voldner, Eva C.; Shannon, Jack D. (1992): Deposition of S and NO_x Nitrogen to the Great Lakes Estimated With A Regional Deposition Model. Environmental Science and Technology (26 May), 970-978. [ATMOSPHERE, GREAT LAKES, POLLUTION, WATER COLUMN]
- Wall, Gregory J. et. al. (1989): Pollution Control in the Great Lakes Basin: an International Effort J Soil & Water Conservation 44, 12-15. [EUTROPHICATION, GREAT LAKES, MANAGEMENT, PHOSPHORUS, POLLUTION, RESTORATION, TOXICS, WATER COLUMN]
- Warwick, W.F.; Casey, C.A. (1982): Sampling Chironomid Communities in Lakes National Water Resources Institute (NWRI) Publication No. W&NR 82-02, NHRI., Saskatoon, Waskatchewan. [BENTHOS, WATER COLUMN]
- The Water Quality Board (1991): Cleaning up in the future. In: Cleaning up Our Great Lakes. A Report From the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 34-36. [GREAT LAKES, PROGRAM, RESTORATION, TRENDS]
- The Water Quality Board (1991): Current Approach to Cleaning up the Lakes. In: Cleaning up Our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 31-33. [GREAT LAKES, PROGRAM, RESTORATION]
- The Water Quality Board (1991): Exposure to and Effects of Toxic Substances. In: Cleaning Up Our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 22-26. [AVIAN SPECIES, DEFORMITIES, FISH, GREAT LAKES, HUMAN HEALTH, MAMMALIAN SPECIES, TOXICS, TRENDS, WILDLIFE]
- The Water Quality Board (1991): Long term goals. In: Cleaning up our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 45. [ECOSYSTEM, GREAT LAKES, HUMAN USE, TRENDS]
- The Water Quality Board (1991): Polluted areas. In: Cleaning up our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 21. [GREAT LAKES, POLLUTION, TOXICS, WATER COLUMN]

The Water Quality Board (1991): Sources of pollution. In: Cleaning up our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 17-20. [ATMOSPHERE, GREAT LAKES, LOADINGS, POLLUTION, SEDIMENT, TOXICS, WASTE DISPOSAL, WATER COLUMN]

The Water Quality Board (1991): Toxic Substances in the Great Lakes. In: Cleaning up our Great Lakes. A Report from the Water Quality Board to the international Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 14-16. [DDT, DIOXIN, GREAT LAKES, LEAD, MERCURY; ORGANOCHLORINE, PCB, PESTICIDE, TOXICS, WATER COLUMN]

The Water Quality Board (1991): Trends in Pollution Discharges and Levels. In: Cleaning up our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 29-30. [GREAT LAKES, LOADINGS, PCB, POLLUTION, TOXICS, TRENDS, WATER COLUMN]

The Water Quality Board (1991): What has been done to clean up the lakes. In: Cleaning up our Great Lakes. A Report from the Water Quality Board to the International Joint Commission on Toxic Substances in The Great Lakes Basin Ecosystem. 27-28. [GREAT LAKES, MANAGEMENT, POLLUTION]

Water Quality and Pollution Control in Michigan: 1990 Report. Michigan 305(b) Report: Volume 11. Lansing, MI. 274 Pages. [WATER QUALITY, LAKE MICHIGAN, GREAT LAKES, TOXICS, RIVERS, SEDIMENTS, AQUATIC LIFE, FISH, HEALTH]

Wells, L;McLin, A.L. (1973): Lake Michigan: Man's Effects on Native Fish Stocks and Other Biota.Great Lakes Fish. Comm. Tech. Rep. 20., Ann Arbor, MI.[FISH, LAKE MICHIGAN]

Whyte, Robert S.; Hartig, John H.; Hopkins, Gordon J. (1990): Decreasing Chloride Trends Observed at Lake Erie Municipal Water Intakes. Journal of Great Lakes Research 16, 233-[LAKE ERIE, LOADINGS, POLLUTION, TRENDS, WATER COLUMN]

Wisconsin Department of Natural Resources (1990): Wisconsin Water Quality Assessment Report to Congress 1990. Section 305(b) Report. 172 pages.[EUTROPHICATION, EXOTIC SPECIES, FISH, HABITAT, HUMAN HEALTH, WATER COLUMN, WETLAND]

Wisconsin Department of Natural Resources (1992): Wisconsin Water Quality Assessment Report to Congress 1992. 244 Pages. (PUBL-WR254-92-REV) [GREAT LAKES, FISH, STREAMS, RIVERS, TOXICS, HUMAN HEALTH, SEDIMENTS, WATER QUALITY, AQUATIC LIFE, TOXICITY TESTING]

2.0 METHODS

2.1 Field Collection Methods

2.1.1 *Benthos*

Benthos samples were collected from the bottom of Lake Michigan at all sampling stations for benthic community analyses and substrate characterization (particle size category and distribution). Samples were collected with a standard Eckman Dredge. Five replicate samples were collected for benthic community analysis and three replicate samples were collected for sediment characterization. When removed from the water, each Eckman Dredge sample was washed into a bucket, then transferred to a 1.0 L wide-mouth plastic sample container. Samples for benthic community analysis were flooded with clean water then preserved with 3% formalin solution prior to shipment for laboratory preparation and analysis. Samples reserved for sediment characterization were not preserved prior to laboratory analysis.

2.1.2 *Phytoplankton*

Phytoplankton samples were collected at each sample location for taxonomic identification and community structure analysis. Four replicate 1.0 L grab samples were collected from composite lake water samples also used for water chemistry samples and chlorophyll-a samples. Lake Michigan water was collected from the surface to approximately one foot above the bottom by a 12 volt battery-powered submersible pump, and composited into a five gallon bucket. Phytoplankton grab samples, chlorophyll-a samples, water chemistry samples, and field determined water quality parameters were obtained from the composite water sample. Phytoplankton samples were preserved with 2.0 to 3.0 mL Lugol's solution prior to shipment and preparation for analysis. Net phytoplankton samples were also obtained concurrent with zooplankton sample collection utilizing an 80 μ m mesh plankton net. Methods are further discussed under the zooplankton section.

2.1.3 Zooplankton

Zooplankton samples were obtained at sites S120, S340, S650 and S1000 by passing a large volume of water through an 80 μ m mesh net of 0.5 meters diameter fitted with a revolving/count flowmeter calibrated for the net size. Prior to deployment of each tow, the beginning number of the flowmeter was recorded. The volume of water sampled was determined by the revolutions and counts attained by the flowmeter during the sample period. The sample net was towed by a motorboat driven at slow speed in circles around each site marker. The net was positioned to the inside of the circle with sufficient distance behind the wash and wake of the boat to intercept undisturbed waters. All zooplankton samples were a composite of material collected from the water column for a timed towing period of three minutes (approximately 6 circles around each site marker). After retrieval of the net, the final flowmeter number was recorded, the net contents were rinsed down to the plankton bucket attached to the bottom of the net, the bucket was completely rinsed into a 500 mL labelled plastic sample container, and the sample was preserved with 3 to 4 mL of Lugol's solution prior to shipment and preparation for analysis.

2.1.4 Ichthyoplankton

Ichthyoplankton (larval fish) samples were collected from S120, S340, S650, and S1000 by using a 300 μ m mesh net towed behind the boat. The ichthyoplankton net was fitted with a revolving/count flowmeter calibrated for the diameter of the net (0.5 meters), and procedures for towing were identical to zooplankton collection methods. All ichthyoplankton samples were collected for a tow period of ten minutes. Upon retrieval of the sampling net, the final number from the flowmeter was recorded, all net contents were rinsed down into the plankton bucket attached to the bottom of the net, and the plankton bucket was completely rinsed into a 500 mL labelled plastic sample container preserved with 3% (vol:vol) formalin solution prior to shipment.

2.1.5 Periphyton

Periphyton growth on site buoy markers (float periphyton) from S120, S340, S650, and S1000 was quantitatively sampled from a known surface area of the buoys. The marker buoys at each

sampling location also served as an artificial substrate for periphyton colonization. SCUBA divers removed the floats from the water to assure minimal disturbance of colonized algae. On board the field boat, several small pieces of buoy approximately 2 to 3 square inches were cut from each marker with a razor blade and placed in Whirl-Pak bags. Deionized water and 1.0 mL of Lugol's solution was added to each sample bag to keep the periphyton wet and preserved until preparation for analysis.

Shore periphyton recognized as the matted filamentous green growth attached to rocks and surfaces along the shoreline. This shore algae is a complex of macro-algae such as the filamentous *Cladophora glomerata*, *Spyrogyra* sp. *Oedogonium* sp. common to most waters. Of recent interest in the Laurentian Great Lakes is the presence and distribution of the red algae (Rhodophyta) *Bangia atropurpurea* in the shore periphyton. Most filamentous algae found along the shoreline environment support an attached complex community of micro-algae including other smaller filamentous forms, diatoms, and other cellular and colonial algae. *Bangia atropurpurea* is not an algae that supports an associated attached (epiphyte) assemblage. As a result, when *Bangia atropurpurea* becomes a significant component of the macro-algae, there is typically a concurrent decrease in the epiphyte community and biomass. According to Lowe *et al* (1982) the lack of epiphyte growth on *Bangia atropurpurea* is a result of the mucilaginous nature of the readily soluble outer cell wall (sulphated galactans) that prevents attachment. This reduction in epiphytic growth would typically reduce biomass and total community diversity. This bioassay samples the shore algae to determine the extent of growth of *Bangia atropurpurea* within and outside the Amoco Cove area.

Shore periphyton attached to rocks situated along the Amoco Cove and Whihala Beach shoreline was qualitatively sampled. Filamentous algae from within the exposed splash zone above the water line, to a depth of approximately 0.5 meters were sampled and composited. Algae were removed from the rock substrates by scraping with a brush, collection with an aspirator, or a combination of both methods. Algae samples were stored in a 250 mL plastic sample container and immediately preserved with Lugol's solution prior to preparation for analysis.

Periphyton associated with the shifting sand bottom substrates was collected to determine if a substantial epipsammon (attached to sand) community existed. Replicate samples of the sandy bottom substrates were collected in 28.2 cm², 118 mL plastic containers with snap-on tops by SCUBA divers. Collection protocols were designed to sample epipsammon as well as refugee algae dislodged from other sites and sources. Collection procedures involved the following steps:

- 1) carefully place the inverted plastic container into the sediments trapping the sand (bottom) and water interface,
- 2) slide a large metal spatula through the surrounding sediments and onto the inverted container to trap the sample,
- 3) remove the container and spatula from the bottom, turn upright slowly without removing spatula seal and let settle for several seconds,
- 4) slowly slide off spatula and snap on sample container top, return to surface.

Four replicate samples of epipsammon algae that is associated with the sandy bottom substrates were collected at each study site. Upon retrieval at the surface, all samples were preserved with 2-3 mL of Lugol's algal preservative and prepared for shipment.

2.1.6 Algal Bioassay

Ambient Lake Michigan waters and undiluted effluent samples were collected on July 25 for use as test conditions for an algal bioassay. Depth composited Lake Michigan water samples were collected in the field at each of S120, S340, and S3500 as described for phytoplankton sample collections. Samples of undiluted effluent were obtained from the Amoco Whiting site prior to the discharge pipe as a composite sample using a permanent automated sampler operated by Amoco personnel. All water and effluent samples were immediately stored on ice and shipped overnight to the ADVENT laboratory.

2.1.7 Water Chemistry

Water chemistry samples for laboratory analysis were collected in the field as a composite of waters from the surface to approximately 1.0 foot above the bottom with a submersible battery-powered pump as described for phytoplankton composite sample collection. Samples for laboratory analysis were transferred to pre-labelled sample bottles by a dipping container to prevent contamination of composited site water. All water chemistry samples were immediately treated according to EPA preservation guidelines and stored on ice until received by the analytical laboratory.

2.1.8 In-situ Water Quality

Water quality parameters that typically change immediately following collection were determined in the field for the composite sample and *in situ* at depth in Lake Michigan as appropriate. The field water quality parameters were determined with the aid of a calibrated Hydrolab Surveyor 3 multi-parameter transmitter and probe. Water quality parameters and physio-chemical parameters recorded *in situ* included the following:

- Total Alkalinity as mg/L CaCO_3 . Determined by titration with weak sulfuric acid to Bromocresol green-methyl red pH 4.5 endpoint.
- pH as standard pH units. Measured by Hydrolab Surveyor 3.
- Dissolved Oxygen in mg/L. Measured by Hydrolab Surveyor 3.
- Specific Conductance (Conductivity) in $\mu\text{mhos/cm}$. Measured by Hydrolab Surveyor 3.
- Temperature in $^{\circ}\text{C}$. Measured by Hydrolab Surveyor 3.
- Depth measured in meters by Hydrolab Surveyor 3.
- Secchi disk depth measured with an 8" standard pattern Secchi disk.

The Hydrolab Surveyor 3 was used to measure temperature, conductivity, dissolved oxygen, pH, and depth at relevant intervals of 1.0 ft from the surface to just above the bottom to determine depth-related water quality profiles.

2.1.9 Effluent Water Quality

A Hydrolab Datasonde 3 datalogger was deployed directly in the Outfall 001 header box. The stand-alone unit was programmed to collect hourly readings of temperature, conductivity, and pH for the two month period of May 12, 1994 to July 12, 1994.

2.2 Laboratory Methods

2.2.1 Benthos

Benthos samples were analyzed by Mr. Henry Zimmermann, a benthic invertebrate entomologist with several years experience in the Great Lakes including southern Lake Michigan. Mr. Zimmermann processed the benthic samples using standard EPA techniques (EPA 1989). All samples were sieved with a 250 μ m screen, then organisms were sorted and identified from the remaining debris from a white enamel pan with the aid of Bausch and Lomb zoom dissecting scope (1-30 X total magnification). A general scan of the sieved sediments was conducted to check for unaccounted specimens. Head capsules of Chironomid specimens were mounted on microscope slides and viewed with a Leitz compound microscope at 100-400 X total magnification for proper identification.

2.2.2 Phytoplankton

The 1.0 L phytoplankton grab samples were checked for sufficient preservation and allowed to completely settle for a minimum of 24 hours following receipt at the AATA laboratory. Approximately 900 mL of water was removed without disturbing the settled material by using a siphon constructed of aquarium hose in order to concentrate the settled phytoplankton into a sample volume of 100 mL. Based on an evaluation of the amount of material settled, each sample was then completely transferred to a 150 mL beaker, covered with a watch glass and allowed to resettle for a period of 24 hours to further concentrate the samples. Final

concentration of the phytoplankton sample to a volume of 20 mL was performed by removal of remaining excess water by siphon.

Concentrated phytoplankton samples were enumerated and identified to the lowest practical taxonomic level at 600 X total magnification using a Palmer-Maloney plankton counting chamber (volume 0.1 mL). Diatoms typically cannot be successfully identified at 600 X total magnification, and were only enumerated for density estimation purposes. Following evaluation of the concentrated phytoplankton sample in three separate preparations of the Palmer-Maloney counting chamber, the total volume of the concentrated sample was measured in a 25 mL graduated cylinder and allowed to settle for a minimum of 24 hours. Following the final settling period, settled material from all site replicates was combined and used to prepare a permanent diatom slide for diatom taxonomic identification and relative percent abundance determination for the sample site. A minimum of 600 diatom frustules were identified from each site to characterize the diatom portion of the phytoplankton community.

2.2.3 Zooplankton

Zooplankton net samples were evaluated by Dr. Edward Reed, formerly of Colorado State University (Professor Emeritus, retired). Zooplankton net samples were completely mixed and split for storage as backup phytoplankton and zooplankton samples for species confirmation. Zooplankton samples used for evaluation were subsampled for observation because of the high number of organisms in the split sample. All organisms from subsamples were sorted and separated from debris then identified and enumerated with the aid of a dissecting microscope ranging from 10 to 130 X total magnification. Identification was augmented by a compound microscope when higher total magnification was required.

2.2.4 Ichthyoplankton

Ichthyoplankton samples were delivered to the Colorado State University Larval Fish Laboratory for analysis. Identification and enumeration of ichthyoplankton samples were conducted using accepted methods and protocols consistent with the standard operational procedures of the Larval Fish Laboratory.

2.2.5 Periphyton

Marker Buoy (Float) Periphyton. Periphyton collected from the marker buoy material was removed by brushing and washing the sample into a 60 mL beaker. Each replicate sample was kept separate and surface areas of all cut pieces of buoy material were measured to determine the total area scraped per replicate sample. The contents of the beaker were measured with a graduated cylinder to determine dilution and concentration factors. Three types of counts were performed to document the taxonomic makeup and abundance data for each sample. First, a 5.0 mL aliquot of thoroughly mixed sample was used to prepare permanent diatom microscope slides for species identification and enumeration at 1,000 to 12,000 X total magnification.

A second evaluation was conducted to determine the density of the diatom portion of the float periphyton. Enumeration of diatom algae cells was conducted with the aid of a hemacytometer (Improved Neubauer ruling) counting chamber at 400 X total magnification. Ten suspensions of completely mixed sample were individually introduced to the hemacytometer (0.0001 mL volume) and all diatoms enumerated within the two grids of the chamber for a total of 20 counts. Diatom density estimates were determined from the volume of the sample, the surface area of the marker buoys samples, and the volume and mean number of diatoms enumerated in the hemacytometer.

A third evaluation was conducted to account for the non-diatom portion of the periphyton and utilized a Palmer-Maloney counting chamber (volume 0.10 mL). A suspension of thoroughly mixed sample was introduced into the Palmer-Maloney counting chamber and non-diatom algae were analyzed at 400 X total magnification for identification and enumeration data. A minimum of 1.5 Palmer Maloney counting chamber values were analyzed per sample. When one half of a Palmer-Maloney counting chamber was analyzed and no new species were encountered, a sufficient sample was believed to be analyzed. Density estimates for the non-diatom algae were determined from the volume of the sample, the surface area of the marker buoy samples, and the volume and enumeration data from the Palmer-Maloney counting chamber analysis. Float periphyton density estimates included diatom periphyton density plus non-diatom periphyton density values.

Shore Periphyton. Matted filamentous periphyton sample replicates collected from rock substrates near Outfall 001 and rock substrates at Whihala Beach ("reference site" located west of the Amoco Facility) were used to determine if gross differences existed in this growth form between the two sites. Prior to viewing with the aid of a microscopic, each sample of scraped algae was individually spread out in a large tray to untangle the algal mat and visually detect the presence of different algal forms by color, size, etc. Microscopic analysis of the shore periphyton included a minimum of 5 large (0.5 area of microscope slide) wet mount preparations of algae, including evaluation of all algal material visually different or unique. Taxa were identified at 400 X total magnification and categorized by one of the three groups for abundance (5=dominant, 3=common, 1=rare). Categories were used in lieu of other quantitative measures because of the inaccuracies and difficulties of sampling and counting a thick filamentous algal mat. The epiphytic algal community attached and associated with the tangled filamentous mat was heavily dominated by a few characteristic diatoms. Separate sample processing for diatom analysis and taxonomy was considered unnecessary and therefore not conducted.

Epipsammon Periphyton. Samples of epipsammon (sand associated) periphyton collected by SCUBA divers from the shifting fine sand substrates required a series of sample treatment steps to separate the algae from the substrates. The procedure used to isolate epipsammon algae from the sandy substrates is listed below:

- 1) Completely irrigate and suspend the sediment with deionized water within the confines of the sample container used in the field.
- 2) Mix the sediment by hand for a period of 10-15 seconds using a teflon laboratory spatula or similar tool to dislodge algae from the sand grains.
- 3) Allow the suspended sediments to settle for a period of approximately 1 - 2 minutes depending upon silt and clay composition of sediments.
- 4) Remove, and retain supernatant water with large bore pipette and store in a 60 mL glass vial or similar suitable container. Minimize collection of sediment material.

- 5) Concentrate the collected algae by allowing the suspension in the glass vial to settle for a period of 12 hours followed by removal of excess water. This step also allows for more supernatant to be added to the sample vial in subsequent steps.
- 6) Repeat steps 1 - 5 above until microscopic evaluation of the last supernatant contains ten or fewer algal cells per field of view (100 X total magnification).

The above procedure removed, isolated and concentrated algae associated with the epipsammon substrate collections.

Analysis of the algae isolated from the epipsammon included initial observation at 360 X total magnification to identify clumps and tangles of macro-algae (primarily filamentous) in a Sedwick-Rafter algal counting chamber (volume 1.0 mL). The initial observations indicated that the epipsammon community consisted of diatoms which were numerically dominated by a few taxa. Subsequently, permanent diatom microscope slides were prepared for identification and enumeration. Due to the dominance of a few taxa, combined with an overall low richness for all sites, a shortened count of 100 frustules for each station was performed at 1000 X total magnification.

2.2.6 Algal Bioassay

Algal bioassay procedures were patterned after the *Selenastrum capricornutum* Printz Algal Assay Bottle Test (USEPA 1978). Two algal species were tested: *Selenastrum capricornutum* and *Scenedesmus quadricauda*. These algae are commonly used in laboratory test procedures. Species of *Scenedesmus* have been found among the phytoplankton in southern Lake Michigan. The algal cultures were obtained from commercial suppliers.

The algal bioassay was designed to test two questions: first, was there any difference in samples taken from outside the effluent dispersion zone from samples taken inside the effluent dispersion zone; and second, would there be any differences expected at dilutions of 20:1 or 40:1 (roughly the dispersion at two sample sites within the effluent dispersion zone). To assure that the test

met quality expectations, two laboratory controls were tested, one using filtered algal growth media and one using unfiltered algal growth media: healthy growth in these controls would indicate that the test was conducted properly and the algae were capable of normal growth. Both algal species were tested in triplicate under all conditions, except that only *Selenastrum* was used in the unfiltered laboratory control tests.

The bioassay design permitted several statistical comparisons, shown in Table 2-1.

Table 2-1. Algal Bioassay Design

Algal Bioassay Design Statistical Comparisons						
Treatment Group		Treatment Comparison				
		A	B	C	D	E
A	S3500 Sample: Lake Michigan reference water outside effluent dispersion zone		1	1	2	3
B	S340 Sample: Estimated 40:1 dilution inside effluent dispersion zone	1				3
C	S120 Sample: Estimated 20:1 dilution inside effluent dispersion zone	1			2	
D	95% S3500 water and 5% Effluent (20:1 dilution)	2		2		
E	97.5% S3500 water and 2.5% Effluent (40:1 dilution)	3	3			
F	Lab Controls: Filtered and unfiltered algal growth medium					

- 1 This comparison evaluates "outside" vs. "inside" effluent dispersion zone.
- 2 This comparison evaluates impact of a 20:1 treated effluent dilution.
- 3 This comparison evaluates impact of a 40:1 treated effluent dilution.

The tests were initiated by inoculating algal cells from rapidly growing (log-phase) cultures into each tests flask at an initial density of 2,000 cells/mL. The test lasted 8 days. The general experimental conditions were:

- Tests were conducted in temperature-controlled room at 25°C under continuous light.
- Lighting was by cool white fluorescent light and reached an intensity of 360 to 440 foot candles at the top and bottom of each flask. Light intensity was measured with a General Electric Model 214 hand-held light meter.
- Each 2000 mL Erlenmeyer flask contained 400 mL of test solution. Flasks were covered with aluminum foil. A small hole in each foil cap permitted gas exchange.
- Flasks were randomly placed on shelves to minimize potential effects of subtle differences due to shelf position. Positions were changed daily, again by random assignment, to further minimize any position effects.

Algal growth was measured by cell counts during the test and by algal dry weight and chlorophyll-a measurement at the conclusion of the test. Chemical parameters were measured on excess media from each treatment group, including analyses for nutrients, major ions and general parameters.

Cell counts were done using a hemacytometer counting slide. A few mL of shaken test solution was withdrawn by pipette and loaded onto one side of the counting slide. A second sample was taken from the same flask and loaded onto the other side. Microscopic cell counts were made and the average of the two sides was recorded as the value for that replicate.

At the end of the 8 day test, 300 mL of each test solution was filtered through a pre-weighed glass fiber filter with 0.7 micron pore size. The algae retained on the filter were dried at 104°C and weighed, giving an algal dry weight. Samples of rapidly-growing algal culture were similarly filtered and weighed to estimate the initial dry weight of the original algal inoculum. The net growth of algae was determined as the difference between day 8 biomass and the initial biomass, and expressed as dry weight of algae per liter.

The filtrate from the 300 mL sample was mixed with other replicates for each treatment and used for chemical analysis. The remaining test sample (approximately 100 mL) was used to measure chlorophyll-a. A measured volume was filtered onto 0.7 micron glass fiber filters, then

the filter and retained algae were sonicated to facilitate extraction on chlorophyll-a with 90% acetone. Chlorophyll-a was measured fluorometrically and expressed as mg chlorophyll-a per liter.

2.3 Analytical and Statistical Methods

Numerical count data associated with species identifications were used to determine the following community structural parameters and characteristics as presented by Ludwig and Reynolds (1988).

- **Richness:** the total number of different taxa,
Hill's N1 value as the number of dominant taxa.
- **Diversity:** Shannon-Weiner Diversity Index,
Simpson's Diversity Index,
Hill's N1 value as a percent of richness.
- **Evenness:** Hill's E5 value.

Richness is a measure of the number of different taxa in a sample collection. Richness values typically represent species level identification, and the values play an important role in determination of other community metrics such as density. Hill's N1 value is the number of abundant species in the sample (e.g. the number of species that are not rare), where each species is weighted by its abundance. Hill's N1 is determined using the Shannon-Weiner Diversity (see discussion below) value as:

$$N1 = e^{H'}$$

where: H' is the Shannon-Weiner Diversity value for the community, and
 $e = 2.7183$

Hill's N1 is an appealing representation of the dominant species in the community because of the easy interpretation with respect to structure and diversity for persons unfamiliar with

diversity indices. Hill's N1 value when compared to total richness gives an intuitive evaluation of how many species in the community may be considered ecologically functional. For convenience, Hill's N1 may be presented as a percent of total richness.

Diversity is a measure of how the abundance is distributed among the different number of taxa (richness). A community with a richness of 20 different taxa, each of which are represented by 15 specimens (total of 300 organisms) will have a much different diversity than 20 different taxa represented by five specimens for each of 18 taxa and 105 specimens for two taxa (total of 300 organisms). Measures of diversity were represented by the Shannon-Weiner Diversity Index (H'); Simpson's Diversity Index (λ); and Hill's N1 value as discussed above. Both the Shannon-Weiner Diversity Index and Simpson's Diversity Index have been used extensively as a descriptive community parameters in ecological studies. The Shannon-Weiner Index considers the abundance of each species in the community and index values increase in magnitude as community diversity increases. A community with one species, regardless of abundance has a Shannon-Weiner Diversity of zero. Maximum values of H' typically range from 3.8 to 4.5 for well developed benthic communities in running waters, or from complex substrates within lakes, and well developed complex planktonic communities.

The Shannon-Weiner Diversity Index values are determined by:

$$\text{Shannon-Weiner Diversity Index: } H' = - \sum_{n=1}^s (p_i \ln p_i)$$

where: p_i = is the proportional abundance of the i th species, and
 s = the total number of species in the community.

The Shannon-Weiner Diversity Index value is used to determine Hill's N1 number of abundant taxa as discussed above.

Simpson's Diversity Index (λ) is a widely used measure of diversity which ranges from zero (for maximum diversity) to 1.0 (for no diversity). Maximum diversity (zero index value) is attained when all species present are of equal abundance. Simpson's Diversity Index is determined by:

$$\text{Simpson's Diversity Index: } (\lambda) = \sum_{i=1}^s \frac{n_i(n_i - 1)}{n(n - 1)} ;$$

where: n_i = the number of individuals of the i th species,
 n = the total number of individuals in the sample, and
 s = the total number of species in the sample.

Evenness has recently become a popular measure of the distribution of species abundance among the total number of taxa much like diversity. However, evenness is typically described as a measure of how relative abundances diverge away from equal abundance among all members of the community. Evenness measured by Hill's modified ratio (E5) is unaffected by species richness, and therefore preferable over other evenness metrics in cases where species richness is expected to be limited. Hill's modified ratio is intuitively easy to understand as the value approaches zero as a single species becomes more and more dominant in the community. Determination of Hill's modified ratio as an evenness metric utilizes Simpson's Diversity Index value, and the Shannon-Weiner Diversity Index value (in the form of Hill's N1 number of abundant taxa). Evenness as measured by Hill's modified ratio is determined by:

$$\text{Hill's modified ratio Evenness: } E5 = \frac{(1/\lambda) - 1}{e^{H'} - 1}$$

where: $e^{H'}$ = Hill's N1 number of abundant taxa,
 λ = Simpson's Diversity Index value.

Complete discussions on the calculations and use of Hill's N1, Simpson's Diversity Index, Shannon-Weiner Diversity Index, and Hill's modified ratio evenness (E5), are presented in Ludwig and Reynolds (1988). Richness, diversity, and evenness community structure parameters were determined with statistical software programs for the IBM provided by Ludwig and Reynolds (1988).

Community structure parameters were used to evaluate biological collections within the dispersion zone and outside the dispersion zone. Sites situated inside the dispersion zone included S120 and S340 as determined by *in-situ* water quality screening and previous dye dispersion studies. Sites situated outside the dispersion zone included S650, S1000, S2000 and S3500. Community structure parameters for all replicate samples inside and outside the dispersion zone were subjected to Analysis of Variance (ANOVA) procedures to indicate if statistical differences existed in mean parameter values inside and outside of the dispersion zone.

Comparisons among communities using structural characteristics is typically sufficient to identify differences between biological samples. However, community structure characteristics are based upon the distribution of organisms among different taxa. Although not common, two communities with nearly identical density and richness would show comparable diversity values but may contain no species in common. An evaluation of the taxonomic makeup of any community is valuable with respect to insight into the ecology and function of the community.

Taxonomic comparisons between the biological collections were performed using the Bray-Curtis Percent Dissimilarity Index (PD). The Bray-Curtis Percent Dissimilarity Index is but one of many similarity indices available, but is favored over presence-or-absence based indices because abundance of each common species is considered in the comparison. The PD is calculated by first:

Calculating Percent Similarity (PS):

$$PS_{jk} = \frac{2W}{A + B} (100)$$

where,

$$W = \sum_{i=1}^s [\min(X_{ij}, X_{ik})] ; \quad A = \sum_{i=1}^s X_{ij} ;$$

and,

$$B = \sum_{i=1}^s X_{ik} ;$$

and,

X_{ij} represents the abundance of the i th species in the j th community,
 X_{ik} represents the abundance of the i th species in the k th community.

A comparison of any two communities with identical species and abundance distributions would result in a percent similarity (PS) of 100%. Bray-Curtis Percent Dissimilarity (PD) is computed by:

$$PD = 100 - PS$$

The Bray-Curtis expression as PD is a useful expression as a distance complement of the PS which may be used in ecological distance based metrics and analyses. The Bray-Curtis PD was used to determine the magnitude of differences in taxonomy and the distribution of taxa among the sample sites. Determination of the PD value was done using the specific statistical software programs for the IBM computer as discussed in Ludwig and Reynolds (1988). Standard descriptive data summary statistics, necessary data transformations and ANOVA procedures were performed using the IBM computer and Microsoft Excel software with added statistical applications.

Statistical procedures and evaluation of algal assay data using *Selenastrum capricornutum* and *Scenedesmus quadricauda* (cell counts, chlorophyll, and dry weight biomass) consisted of analysis of variance (ANOVA) procedures performed to compare all treatments for a given

species. If significant differences were found, the appropriate multiple range tests (Dunnett's procedure for parametric data and Wilcoxon's procedure for nonparametric data) were performed to identify significantly different exposures. To compare a field sample to analogous laboratory-prepared samples, the Student's t-test was performed. Statistical significance was determined at the 95 percent confidence level.